

Appendix: I-CARE Collaborator Group

Investigators, listed by decreasing number of patients enrolled:

Jean-Marc GORNET, Hôpital Saint-Louis, Université Paris Nord, Paris, France; Laurent BEAUGERIE, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Sebastian SHAJI, Hull University Teaching Hospitals, Hull York Medical School - University of Hull, Hull, UK; Laurent PEYRIN-BIROULET, CHRU Nancy, Nancy University, Nancy, France; Jean-Marie REIMUND, Hôpital universitaire de Strasbourg, Université de Strasbourg, Strasbourg, France; Xavier HEBUTERNE, Hôpital L'archet, Nice, France; Aurélien AMIOT, Hôpital Henri Mondor Université Paris Est, Créteil, France; Franco ARMELAO, Ospedale Santa Chiara, Azienda Provinciale per i Servizi Sanitari, Trento, Italy; Pierre BLANC, CHU Montpellier, Université de Montpellier, Montpellier, France; Claudio PAPI, Ospedale San Filippo Neri, Roma, Italy; Guillaume PINETON DE CHAMBRUN, Hôpital Saint-Eloi, Université de Montpellier, Montpellier, France; Xavier ROBLIN, CHU Saint-Etienne, Saint-Etienne Université, Saint-Etienne, France; Konstantinos KARMIRIS, Venizeleio General Hospital, Heraklion, Greece; Sohail SHARIQ, Ealing Hospital, Imperial college london, London, UK; Nikolaos VIAZIS, Evangelismos General Hospital, Athens, Greece; Jimmy LIMDI, Northern Care Alliance NHS Foundation Trust, University of Manchester, Manchester, UK; Piotr EDER, H. Święcicki University Hospital, Poznan University of Medical Sciences, Poznan, Poland; Georgios MICHALOPOULOS, Tzaneion General Hospital, Pireaus, Greece; Andrew BELL, Weston General Hospital, Weston-Super-Mare, UK; Livia BIANCONE, Policlinico Tor Vergata, Università di Roma Tor Vergata, Roma, Italy; Marie DEWITTE, CHU Rennes - Hôpital Pontchaillou, Université Rennes 1, Rennes, France; Zia MAZHAR, Basildon and Thurrock University Hospitals NHS Trust, Basildon, UK; Denis FRANCHIMONT, Erasmus Hospital, VUB -ULB, Brussels, Belgium; Stephane NANCEY, Hôpital Lyon-Sud, Université

Lyon 1, Lyon, France; Gilles MACAIGNE, CH Marne la vallée, Jossigny, France; Maria Beatrice PRINCIPI, Policlinico di Bari, Università di Bari, Bari, Italy; Mathurin FUMERY, CHU Amiens, Université de Picardie Jules Verne, Amiens, France; Gareth PARKES, St Bartholomew's Hospital, Queen Mary University of London, London, UK; Jean-Christophe VALATS, CHU Montpellier, Université de Montpellier, Montpellier, France;; Glen DOHERTY, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Guillaume BOUGUEN, CHU Rennes - Hôpital Pontchaillou, Université Rennes 1, Rennes, France; Tamás MOLNAR, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary;; Hersin TSAI, Hull University Teaching Hospitals, Hull York Medical School - University of Hull, Hull, UK; Mohsin GANGI, Stepping Hill Hospital, Stockport, UK; Natalia PEDERSEN, Slagelse Hospital, University of Copenhagen, Slagelse, Denmark; Frédéric HELUWAERT, CH Annecy Genevois, Annecy, France; Richard SHENDEREY, Airedale General Hospital, Leeds University School of Medicine, Airedale, UK; Sebastian ZEISSIG, , University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany; Jeffrey BUTTERWORTH, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Fabiana CASTIGLIONE, Policlinico Federico II, Università Federico II, Napoli, Italy; Lynsey CORLESS, Hull University Teaching Hospitals, Hull York Medical School- University of Hull, Hull, UK; Camille ZALLOT, CHRU Nancy, Université de Nancy, Nancy, France; ; Filip BAERT, AZ Delta, campus Rumbeke, Roeselare, Belgium; Salil SINGH, Royal Bolton Hospital, Bolton, UK; Sunil SONWALKAR, Calderdale Royal Hospital, Halifax, UK; Julie BURTON , Pinderfields Hospital, Wakefield, UK; Elizabeth CLAYTON, Pinderfields Hospital, Wakefield, UK; Jean-François RAHIER, CHU Namur, site Godinne, University of Namur, Namur, Belgium; Deven VANI, Pinderfields Hospital, Wakefield, UK; Renata D'INCA, Ospedale di Padova, Università di Padova, Padova, Italy; Guy BELLAICHE, Hôpital Robert Ballanger, Université de Bobigny, Aulnay-sous-bois,

France; Martine DE VOS, UZ Ghent, Ghent University, Ghent, Belgium; Julien KIRCHGESNER, Hôpital Saint-Antoine, Sorbonne Université , Paris, France; Uri KOPYLOV, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Triana LOBATON, UZ Ghent, Ghent University, Ghent, Belgium; Christophe LOCHER, Hôpital de Meaux, Meaux, France; Gerassimos MANTZARIS, Evangelismos General Hospital, Athens, Greece; George ABOUDA, Hull University Teaching Hospitals, Hull York Medical School - University of Hull, Hull, UK; Katie SMITH, Michael SPRAKES, Pinderfields Hospital, Wakefield, UK; Yeovil District Hospital, Yeovil, UK; Angeliki THEODOROPOULOU, Venizeleio General Hospital, Heraklion, Greece; Emma WESLEY, Musgrove Park Hospital, Taunton, UK; Joëlle BONNET, Hôpital Saint-Louis, Université Paris Nord, Paris, France; David ELPHICK, Chesterfield Royal Hospital, Chesterfield, UK; Cyrielle GILLETTA, Hôpital Rangueil, Université Toulouse 3 , Toulouse, France; John GORDON, Hampshire Hospitals, University of Southampton, Basingstoke, UK; David LAHARIE, CHU Bordeaux, Université de Bordeaux , Pessac, France; Antoine NAKAD, Chwapi Hospital, Tournai, Belgium; Ambrogio ORLANDO, Ospedale V. Cervello, Università di Palermo, Palermo, Italy; Patrick DUBOIS, Kings College Hospital, King's College London, London, UK; Peter HASSELBLATT, Universitätsklinikum Freiburg, Universitätsklinikum Freiburg, Freiburg, Germany; Christophe MICHIELS, CHU Dijon, Dijon, France; Cathryn PRESTON, Bradford Teaching Hospitals, Leeds School of Medicine, Bradford, UK; Anca STAICU, Hull University Teaching Hospitals, Hull York Medical School/ University of Hull, Hull, UK; Lucine VUITTON, CHU de Besançon - UMR, Université de Besançon, Franche-comté, France; Mehdi KAASSIS, Centre Hospitalier de Cholet, Cholet, France; Ally SPEIGHT, Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Deb GHOSH, Princess Alexandra Hospital, Harlow, UK; Mark LÖWENBERG, Amsterdam University Medical Center, location AMC, Amsterdam, The Netherlands; Nicolas MATHIEU, CHU Grenoble,

Université Grenoble Alpes , Grenoble, France; Anne-Laure PELLETIER, Groupe Hospitalier Bichat-Claude Bernard, Paris, France; Anne PHILLIPS, York Hospital, Hull York Medical School, York, UK; Fernando MAGRO, Centro Hospitalar Universitário de São João, Faculty of Medicine of the University of Porto, Porto, Portugal; Romain ALTWEGG, Hôpital Saint-Eloi, Montpellier, France; Irit AVNI BIRON, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Jonathon LANDY, Watford General Hospital, Watford, UK; Maria NACHURY, CHU Lille, Université Lille, Lille, France; Achuth SHENOY, Colchester General Hospital, Barts and The London School of Medicine and Dentistry Norwich Medical School Anglia Ruskin University, Colchester, UK; Caroline TRANG, CHU Nantes, Université Nantes , Nantes, France; Vered ABITBOL, Hôpital Cochin AP-HP, Université Paris Cité, Paris, France; Georgios BAMIAS, Sotiria General Hospital, National and Kapodistrian University, Athens, Greece; Klaudia FARKAS, Albert Szent-Györgyi Health center, University of Szeged, Szeged, Hungary; Christian MAASER, Klinikum Lüneburg, Klinikum Lüneburg, Lüneburg, Germany; Ariella SHITRIT, Shaare Tzedek Medical Center, Hebrew University, Jerusalem, Israel; Britta SIEGMUND, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Jérôme FILIPPI, Hôpital L'archet, Nice, France; Colm O'MORAIN, Beacon Hospital, UCD, Dublin, Ireland; Henit YANAI, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Laurent COSTES, Centre hospitalier intercommunal de Créteil, Créteil, France; David HOBDAY, Sunderland Royal Hospital, Sunderland, UK; Zoltán SZEPES, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Emma CALABRESE, Policlinico Tor Vergata, Università di Roma Tor Vergata, Roma, Italy; Helen DALLAL, James Cook University Hospital, Durham University, Newcastle University, Middlesbrough, UK; Michael FUNG, NHS Plymouth, Plymouth, UK; Arvind RAMADAS, James Cook University Hospital, Durham University, Newcastle University, Middlesbrough, UK; Bijay BABURAJAN,

Maidstone Hospital, Maidstone, UK; Konrad KOSS, Macclesfield District General Hospital, Macclesfield, UK; Christophe BARBERIS, Hôpital Bagatelle, Talence, France; Anthony BUISSON, CHU Clermont-Ferrand, Inserm U1071, M2iSH, Université Clermont Auvergne, Clermont-Ferrand, France; Morgane AMIL, CHD Vendée, La Roche-sur-Yon, France; Paola BALESTRIERI, Campus Biomedico, Università Campus Biomedico, Roma, Italy; Matthew JOHNSON, Luton and Dunstable University Hospital, University College London, Luton, UK; Maria TZOUVALA, Agios Panteleimon General Hospital, Nikaia-Piraeus, Greece; Stéphanie VIENNOT, CHU Caen, Université de Caen, Caen, France; Andy LI, Worthing Hospital, Worthing, UK; Ferenc NAGY, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Nick THOMPSON, Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Laurent ALRIC, Hôpital Rangueil, Université Toulouse 3, Toulouse, France; Sunil SAMUEL, Nottingham University Hospitals NHS Trust Queen's Medical Centre, Nottingham, UK; Anne BOURRIER, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Elise CHANTELOUP, Hôpital Paris Saint-Joseph, Paris, France; Emilie DEL TEDESCO, CHU Saint-Etienne, Université Saint-Etienne, Saint-Etienne, France; Marcus HARBORD, Chelsea and Westminster Hospital, Imperial college london, London, UK; Alan LOBO, Royal Hallamshire Hospital, University of sheffield, Sheffield, UK; Sally MYERS, Hull University Teaching Hospitals, Hull York Medical School - University of Hull, Hull, UK; Richard POLLOK, St George's Hospital, University of London, London, UK; Tariq AHMAD, Royal Devon and Exeter Hospital, University of Exeter Medical School, Exeter, UK; Rakesh CHAUDHARY, Watford General Hospital, Watford, UK; Christos KARAKOIDAS, Naval Hospital, Athens, Greece; Edouard LOUIS, CHU de Liège, ULiège, Liège, Belgium; Ashraf SOLIMAN, Barnsley General Hospital, University of Sheffield, Barnsley, UK; Carmen STEFANESCU, Hôpital Beaujon, Université Paris, Clichy, France; Georgios THEOCHARIS, University Hospital of Patra, Patra, Greece; Stijn VANDEN BRANDEN, OLV-Aalst, Aalst,

Belgium; Belén BELTRAN , Hospital Universitari La Fe - Instituto de Investigación Sanitaria La Fe (IISLaFe), Polytechnic University of Valencia, Valencia, Spain; Yoram BOUHNİK, Hôpital Beaujon, Université de Paris, Clichy, France; Arnaud BOURREILLE, CHU Nantes, Université de Nantes , Nantes, France; Joana BRANCO, Hospital Prof. Doutor Fernando Fonseca, Amadora, Portugal; Ben COLLEYPRIEST, Royal United Hospitals Bath, Bath, UK; Rami ELIAKIM, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Paul KNIGHT, Wythenshawe Hospital, University of Manchester, Manchester, UK; Aoibhlinn O'TOOLE, Beaumont Hospital, RCSI, Dublin, Ireland; Virginia ROBLES, Hospital Vall d'Hebron, Universidad autonoma de Barcelona, universitat oberta de catalunya, Barcelona, Spain; Konstantinos TRIANTAFYLLOU, Attikon University Hospital, National and Kapodistrian University, Athens, Greece; Marta Maia BOSCA, Hospital Clinico Universitario de Valencia, University of Valencia, Valencia, Spain; Guy LAMBRECHT, AZ Damiaan, Ostend, Belgium; Lucia MARQUEZ MOSQUERA, Hospital del Mar. Barcelona, Pompeu Fabra University, Barcelona, Spain; Simon PANTER, South Tyneside District Hospital, South Shields, UK; Aikaterini PAPPA, University Hospital of Larissa, Larissa, Greece; Marion SIMON, Institut Mutualiste Montsouris, Paris, France; Ganesh SIVAJI, Harrogate District Hospital, Harrogate, UK; Christophe BELLANGER, Kremlin Bicêtre, Université Paris Saclay, Le Kremlin Bicêtre, France; Arthur BELLE, CHRU Nancy, Université de Nancy , Nancy, France; Natalia BORRUEL, Hospital Vall d'Hebron and Centro de Investigación Biomédica en Red de enfermedades Hepáticas y Digestivas (CIBEREHD), Autonomous University of Barcelona, Barcelona, Spain; Laurence EGAN, UCHG, NUIG, Galway, Ireland; Harald PEETERS, AZ St-Lucas, Ghent, Ghent, Belgium; Daniel SHARPSTONE, West Suffolk Hospital, University of Suffolk, Suffolk, UK; Ramesh ARASARADNAM, University Hospital Coventry, University of Warwick, Coventry, UK; José Manuel BENITEZ, Hospital Universitario Reina Sofía, Instituto Maimónides de Investigación

Biomédica de Córdoba (IMIBIC), University of Cordoba, Córdoba, Spain; Jens Frederik DAHLERUP, Aarhus University Hospital, University of Aarhus , Aarhus, Denmark; Olga GIOULEME, Hippokration General Hospital, Aristotle University, Thessaloniki, Greece; Javier P. GISBERT, Hospital Universitario de La Princesa, Instituto de Investigación Sanitaria Princesa (IIS-IP) and Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Autonomous University of Madrid (UAM), Madrid, Spain; Ulf HELWIG, Internistische Praxengemeinschaft Oldenburg, University of Kiel, Germany; Miguel MINGUEZ, Hospital Clinico Universitario de Valencia, University of Valencia, Valencia, Spain; Eftychia TSIRONI, Metaxa Memorial General Hospital, Pireaus, Greece; Angela VARIOLA, Ospedale Sacro Cuore Don Calabria, Negrar di Valpolicella, Italy; Patrick ALLEN, Ulster Hospital, Belfast, UK; Lucille BOIVINEAU, Hôpital Saint-Eloi, Montpellier, France; Andy COLE, Royal Derby Hospitals, Derby, UK; Nina DIB, CHU Angers, Université Angers , Angers, France; Fernando GOMOLLON, Hospital Clínico Universitario "Lozano Blesa", Instituto de Investigación Sanitaria de Aragón (IIS-Aragón), Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), University of Zaragoza, Zaragoza, Spain; Richard JOHNSTON, Torbay Hospital, Torquay, UK; Konstantinos KATSANOS, University Hospital of Ioannina, University of Ioannina, Ioannina, Greece; Nick KENNEDY, Royal Devon and Exeter Hospital University of Exeter Medical School, Exeter, UK; Marianne KISZKA-KANOWITZ, Hvidovre Hospital, University of Copenhagen, Hvidovre, Denmark; Ignacio MARIN-JIMENEZ, Hospital Universitario Gregorio Marañón, IiSGM, Complutense University of Madrid, Madrid, Spain; Pál MIHELLER, 1st Department of Surgery and Interventional Gastroenterology, Semmelweis University, Budapest, Hungary; Pilar NOS, Hospital Universitari La Fe, Instituto de Investigación Sanitaria La Fe (IISLaFe), Universitat Politècnica de Valencia, Valencia, Spain; Othman SARAJ, Scarborough Hospital, Hull York Medical School, Scarborough, UK; Lars

VINTER-JENSEN, Aalborg University Hospital, University of Aalborg, Aalborg, Denmark; Eran ZITTAN, Haemeq Medical Center, Faculty of Medicine Technion, Afula, Israel; Clotilde BAUDRY, Hôpital Saint-Louis, Université Paris Nord, Paris, France; Xavier CALVET, Hospital Universitari Parc Taulí, Autonomous University of Barcelona, Sabadell, Spain; Marie-Christine CAZELLES-BOUDIER, Centre Hospitalier de la Côte Basque, Bayonne, France; Jean-Louis COENEGRACHTS, Jessa Ziekenhuis Hospital, Hasselt, Belgium; Garret CULLEN, St Vincent University Hospital & University College Dublin, Dublin , Ireland; Marco DAPERNO, Ospedale Ordine Mauriziano Umberto I, Università di Torino, Torino, Italy; Anjan DHAR, Darlington Memorial Hospital, Durham, UK; Romain GERARD, CHU Lille, Université de Lille, Lille, France; Christian Lodberg HVAS, Aarhus University Hospital, University of Aarhus , Aarhus, Denmark; Nanna JENSEN, Bispebjerg Hospital, University of Copenhagen, Copenhagen, Denmark; Nitsan MAHARSHAK, Sourasky Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Mark MCALINDON, Royal Hallamshire Hospital, University of sheffield, Sheffield, UK; Simon MCLOUGHLIN, Christchurch Hospital, Christchurch, UK; Miles PARKES , Addenbrookes's Hospital, University of Cambridge, Cambridge, UK; Kamal PATEL, St George's Hospital, University of London, London, UK; Armando PEIXOTO, Centro Hospitalar Universitário de São João, Faculty of Medicine of the University of Porto, Porto, Portugal; Dimitrios POLYMEROS, Attikon University Hospital, Athens, Greece; Francisco PORTELA, Centro Hospitalar Universitário de Coimbra, Coimbra, Portugal; Rodolfo ROCCA, Ospedale Ordine Mauriziano Umberto I, Università di Torino, Torino, Italy; Philippe SEKSIK, Hôpital Saint Antoine, Sorbonne Université, Paris, France; Sreedhar SUBRAMANIAN, Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Ruth TENNENBAUM , Centre Hospitalier Poissy St Germain-en-Laye, St Germain-en-Laye, France; Raja ATREYA, Universitätsklinikum

Erlangen, Universitätsklinikum Erlangen, Erlangen, Germany; Oliver BACHMANN, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover, Hannover, Germany; Arthur BERGER, CHU Angers, Université Angers, Angers, France; Renáta BOR, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Maire BUCKLEY , St Vincent University Hospital & University College Dublin, Dublin, Ireland; Daniel CARPIO, Complejo Hospitalario Universitario de Pontevedra, University of Alcalá de Henares and the University of Granada, Pontevedra, Spain; María CHAPARRO, Hospital Universitario de La Princesa, Instituto de Investigación Sanitaria Princesa (IIS-IP) and Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Autonomous University of Madrid (UAM), Madrid, Spain; Francesco COSTA, Azienda ospedaliero-universitaria pisana, Università di Pisa, Pisa, Italy; Eugeni DOMENECH, Hospital Universitari Germans Trias i Pujol Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Universitat Autònoma Barcelona, Badalona, Spain; Maria ESTEVE, Hospital Universitari Mutua Terrassa. Centro Investigación biomédica en red de enfermedades hepaticas y digestivas, University of Barcelona, Barcelona, Spain; Stephen FOLEY, King's Mill Hospital, University of Nottingham, Sutton-in-Ashfield, UK; Jordi GUARDIOLA, Hospital Universitario de Bellvitge, IDIBELL, Universidad de Barcelona, Hospitalet del Llobregat, Spain; Ioannis KOUTROUBAKIS, University Hospital of Heraklion, University of Crete, Heraklion, Greece; Tanja KUEHBACHER, Medius Kliniken Nürtingen, Medius Kliniken Nürtingen, Nürtingen, Germany; Cécilia LANDMAN, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Alessandro LAVAGNA, Ospedale Ordine Mauriziano Umberto I, Università di Torino, Torino, Italy; Noemí MANCENÍDO, Hospital Universitario Infanta Sofía, European University of Madrid, San Sebastián de los Reyes, Spain; Míriam MAÑOSA, Hospital Universitari Germans Trias i Pujol Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Badalona, Spain;

Maria Dolores MARTÍN-ARRANZ, La Paz University Hospital, Universidad Autónoma de Madrid, Hospital La Paz Institute for Health Research, La Paz Hospital, Madrid, Spain; Laurianne PLASTARAS, CHU Colmar, Colmar, France; Maria Lia SCRIBANO, Ospedale S. Camillo-Forlanini, Roma, Italy; Subhasish SENGUPTA, Our Lady of Lourdes Hospital, RCSI, Drogheda, Ireland; Nils TEICH, Internistische Gemeinschaftspraxis für Verdauungs- und Stoffwechselkrankheiten, Leipzig, Germany; My-Linh TRAN-MINH, Hôpital Saint-Louis, Université Paris Nord , Paris, France; Evanthia ZAMPELI, Alexandra General Hospital, Athens, Greece; Leila AMININEJAD, Erasmus Hospital, VUB-ULB, Brussels, Belgium; Teresa ARROYO, Hospital Clínico Universitario "Lozano Blesa", Instituto de Investigación Sanitaria de Aragón (IIS-Aragón), Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), University of Zaragoza, Zaragoza, Spain; Alain ATTAR, Hôpital Beaujon, UFR de Médecine, Xavier Bichat, Université de Paris, Clichy la garenne, France; Ann-Sofie BACKMAN, Karolinska University Hospital, Inst Med Solna Karolinska Institutet, Stockholm, Sweden; Anita BÁLINT, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; John BECKLY, Royal Cornwall Hospital, University of Exeter Medical School, Truro, UK; Shomron BEN HORIN, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Sónia BERNARDO, Centro Hospitalar Lisboa Norte, Lisboa, Portugal; Ludovic CAILLO, Hôpital universitaire de Nîmes, Université Montpellier-Nîmes, Nîmes, France; Bénédicte CARON, Hôpital Universtaire de Strasbourg, Université de Strasbourg , Strasbourg, France; María Shanika DE SILVA, Russells Hall Hospital, Dudley, UK; Anna FÁBIÁN, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Gionata FIORINO, IRCCS S. Raffaele, Università Vita-Salute San Raffaele, Milano, Italy; Ana GUTIERREZ, Hospital General Universitario Alicante, Centro de Investigación Biomédica en Red de enfermedades Hepáticas y Digestivas - ISABIAL, Instituto de Investigación Sanitaria y Biomédica de Alicante, Alicante, Spain; Adi LAHAT, Sheba

Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Mohamed MASMOUDI, Hôpital de Meaux, Meaux, France; Marco MENDOLARO, Ospedale Ordine Mauriziano Umberto I, Università di Torino, Torino, Italy; Vinciane MULS, CHU Saint-Pierre, VUB -ULB, Brussels, Belgium; Florian POULLENOT, CHU Bordeaux, Université de Bordeaux, Pessac, France; Christopher PROBERT, Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Catherine REENAERS, CHU de Liège, ULiège, Liège, Belgium; Mariann RUTKA, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Zaman SARWARI, Odense University Hospital Svendborg, University of Odense, Svendborg, Denmark; Joanne SAYER, Doncaster Royal Infirmary, Doncaster, UK; Beatriz SICILIA, Hospital Universitario de Burgos, Burgos, Spain; Helena SOUSA, Centro Hospitalar Universitário do Algarve - Portimão, Portimão, Portugal; Catherine VAN KEMSEKE, CHU de Liège, ULiège, Liège, Belgium; Ana VIEIRA, Hospital Garcia Orta, Almada, Portugal; Yamile ZABANA, Hospital Universitari Mútua Terrassa, Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), University of Barcelona, Terrassa, Spain; Marco ASTEGIANO, Ospedale San Giovanni antica sede, AOU Città della Salute e della Scienza di Torino, Torino, Italy; Paul BANIM, James Paget University Hospital, University of Suffolk, Great Yarmouth, UK; Dominik BETTENWORTH, Praxis für Innere Medizin, Münster, Germany; Médina BOUALIT, Hôpital de Valenciennes, Université de Lille, Valenciennes, France; Jacob Broder BRODERSEN, Hospital of Southwest Jutland, University of Southern Denmark, Esbjerg, Denmark; Angeliki CHRISTIDOU, Evangelismos General Hospital, Athens, Greece; Rachel COONEY, Queen Elizabeth Hospital, University of Birmingham, Birmingham, UK; João CORTEZ PINTO, Instituto Português Oncologia - Lisboa Francisco Gentil, Lisboa, Portugal; Marília CRAVO, Hospital Beatriz Ângelo, Loures, Portugal; Anneline CREMER, Erasmus Hospital, VUB-ULB, Brussels, Belgium; Silvio

DANESE, IRCCS S. Raffaele, Università Vita-Salute San Raffaele, Milano, Italy; Antonio DI SABATINO, Policlinico San Matteo, Università di Pavia, Pavia, Italy; Jan FALLINGBORG, Aalborg University Hospital, University of Aalborg, Aalborg, Denmark; Antonio FERRONATO, Ospedale Alto Vicentino AULSS 7 Pedemontana, Santorso, Italy; Esther GARCIA PLANELLA, Hospital de la Santa Creu i Sant Pau, Open University of Catalonia, Barcelona, Spain; Sanjay GUPTA, Croydon University Hospital, St George's, University of London King's College School of Medicine and Dentistry, Croydon, UK; Jonas HALFVARSON, Örebro University Hospital, Faculty of Medicine and Health, Örebro University, Örebro, Sweden; Eran ISRAELI, Wolfson Medical Center, Sackler Medical School Tel Aviv University, Tel Aviv, Israel; Samantha KESTENBAUM, Bristol Royal Infirmary, University of Bristol, Bristol, UK; Lone LARSEN, Aalborg University Hospital, University of Aalborg, Aalborg, Denmark; Elisabeth MACKEN, UZA, Antwerp University, Antwerp, Belgium; Nicoletta MATHOU, Konstantopoulio General Hospital, Athens, Greece; Ágnes MILASSIN, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Joanna POFELSKI, CH Annecy Genevois, Annecy, France; Chiara RICCI, Presidio Spedali Civili di Brescia, Università di Brescia, Brescia, Italy; Francisco RODRIGUEZ-MORANTA, Hospital Universitario de Bellvitge, IDIBELL, Hospitalet del Llobregat, Spain; Martin SCHMIDT-LAUBER, Praxis für Gastroenterologie Oldenburg, Oldenburg, Germany; Ian SHAW, Gloucester Royal Infirmary, Gloucester, UK; Marta SOARES, Centro Hospitalar Universitário de Coimbra, Coimbra, Portugal; Heithem SOLIMAN, Hôpital Louis Mourier, Université de Paris , Paris, France; Christos TRIANTOS, University Hospital of Patra, University of Patras, Patra, Greece; Konstantinos ZOGRAFOS, Athens Medical P. Faliron, Athens, Greece; Anurag AGRAWAL, Doncaster Royal Infirmary, Doncaster, UK; Alessandro ARMUZZI, IBD Center Humanitas Research Hospital , Rozzano Department of Biomedical Sciences Humanitas University, Peive Emanuele, Milan, Italy; Alexandre AUBOURG, CHU

Trousseau Tours, Université de Tours, Tours, France; Manuel BARREIRO-DE ACOSTA, Hospital Clínico Universitario de Santiago de Compostela, University of Santiago de Compostela, Santiago, Spain; Jesús BARRIO, Hospital Universitario Río Hortega, Gerencia Regional de Salud de Castilla y León (SACYL), Valladolid, Spain; Daniel BERGEMALM, Örebro University Hospital, Faculty of Medicine and Health, Örebro University, Örebro, Sweden ; Fernando BERMEJO, Hospital Universitario de Fuenlabrada, University Rey Juan Carlos, Madrid, Spain; Giorgia BODINI, Ospedale San Martino, Università di Genova, Genova, Italy; Johan BOHR, Örebro University Hospital, Faculty of Medicine and Health, Örebro University, Örebro, Sweden; Peter BOSSUYT, Imelda Hospital, Bonheiden, Belgium; Dimitrios CHRISTODOULOU, University Hospital of Ioannina, University of Ioannina, Ioannina, Greece; Christophe CLAESSENS, AZ Turnhout, Turnhout, Belgium; Paul COLLINS, Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Ruth DE FRANCISCO, Hospital Universitario Central de Asturias, Instituto de Investigación Sanitaria del Principado de Asturias (ISPA), University of Oviedo, Oviedo, Spain; Santiago GARCIA, Hospital Universitario Miguel Servet, Zaragoza University, Zaragoza, Spain; Sotirios GEORGOPOULOS, Athens Medical P. Faliron, Athens, Greece; Felix GOUTORBE, Centre Hospitalier de la Côte Basque, Bayonne, France; Chrisostomos KALANTZIS, Army Share Fund Hospital, Athens, Greece; Anastasia KOURIKOU, Hippokration General Hospital, Athens, Greece; Vincent MACE, CHD Vendée, La Roche-sur-Yon, France; Georgia MALAMUT, Hôpital Européen Georges Pompidou, Université de Paris, Paris, France; Paula MINISTRO, Centro Hospitalar Tondela-Viseu, Viseu, Portugal; Isabelle NION LARMURIER, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Elena RICART, Hospital Clínic Barcelona Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD) IDIBAPS, University of Barcelona (UB), Barcelona, Spain; Mélanie SERRERO,

Hôpital Nord Marseille, Université d'Aix-Marseille , Marseille, France; Juliette SHERIDAN, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Petra WEIMERS, North Zealand Hospital, University of Copenhagen, Hillerød, Denmark; Vibeke ANDERSEN, Hospital of Southern Jutland, University of Southern Denmark, Aabenraa, Denmark; Bruno ARROJA, Hospital de Braga, Braga, Portugal; Bernd BOKEMEYER, Gastroenterologische Gemeinschaftspraxis Minden, Minden, Germany; Luis BUJANDA, Hospital Universitario de Donostia. Biodonostia Health Research Institute. Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), University of the Basque Country (UPV/EHU), San Sebastián, Spain; Thibault DEGAND, CHU Dijon, Dijon, France; Carl ERIKSSON, Örebro University Hospital, Faculty of Medicine and Health, Örebro University, Örebro, Sweden; Cécile GARCEAU, CHD Vendée, La Roche-sur-Yon, France; Henning GLERUP, Regional Hospital Silkeborg, University of Aarhus, Silkeborg, Denmark; Idan GOREN, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Lucina JACKSON , Bon Secours Hospital, UCC, Cork, Ireland; Stéphane KOCH, CHU Besançon, Université Besançon, Franche-comté, France; Francisco MESONERO, Hospital Universitario Ramón y Cajal, University of Alcalá de Henares, Madrid, Spain; Ingrid ORDAS, Hospital Clinic of Barcelona, Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBER-EHD), Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona (UB), Barcelona, Spain; Pauline RIVIERE, CHU Bordeaux, Université de Bordeaux, Pessac, France; Simone SAIBENI, Ospedale di Rho, Rho, Italy; João SOARES, Hospital de Braga, Braga, Portugal; Noémie TAVERNIER, Hôpital Tourcoing, Université de Lille, Tourcoing, France; Klaus THEEDE, Hvidovre hospital, University of Copenhagen, Hvidovre, Denmark; Bella UNGAR, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Elke BÄSTLEIN, Magen Darm Zentrum Wiener Platz, Köln, Germany; Antonio GASBARRINI,

Policlinico A. Gemelli, Università Cattolica del Sacro Cuore, Italy; Andreas PROTOPAPAS, AHEPA University Hospital, Thessaloniki, Greece; Wolfgang REINDL, Universitätsmedizin Mannheim, Universitätsmedizin Mannheim, Mannheim, Germany; Fabrizio BOSSA, Fondazione Casa Sollievo della Sofferenza, IRCCS , San Giovanni Rotondo, Italy; Ailsa HART, St Marks Hospital, Imperial college london, London, UK; Franz-Josef HEIL, Fachärztliche Schwerpunktpraxis für Magen-Darm-Erkrankungen, Andernach, Germany; Anthony O'CONNOR, TUH, TCD, Dublin, Ireland; Bas OLDENBURG, UMC Utrecht, Utrecht University, Utrecht, The Netherlands; Luca PASTORELLI, Policlinico San Donato, Università di Milano, San Donato Milanese, Italy; Stephen PATCHETT, Beaumont Hospital, RCSI, Dublin, Ireland; Subramaniam RAMAKRISHNAN, Warrington Hospital, Warrington, UK; John De CAESTECKER, Leicester Royal Infirmary, Leicester University, Leicester, UK; Ana ECHARRI, Hospital ArquiStudy Coordinatorto Marcide, Complejo Hospitalario Universitario de Ferrol, University of Navarra, Ferrol, Spain; David KEVANS, St. James's Hospital, TCD, Dublin, Ireland; Jürgen BÜNING, Schwerpunktpraxis Gastroenterologie, Lübeck, Germany; Rosa COELHO, Centro Hospitalar Universitário de São João, Faculty of Medicine of the University of Porto, Porto, Portugal; Jeroen JANSEN, OLVG oost, Amsterdam, The Netherlands; Benjamin KOSLOWSKI, Shaare Tzedek Medical Center, Hebrew University, Jerusalem, Israel; Christopher WELLS, University Hospital of North Tees, Stockton, UK; Daniel CEBALLOS, Hospital Universitario de Gran Canaria Dr. Negrín, University of Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain; Daan De Konig, Gelre ziekenhuizen, Gelre, The Netherlands; Ingrid KÖNIG, Praxis Hopsital, Bautzen, Germany; Hari PADMANABHAN, Newcross Hospital, Wolverhampton, UK; Timi PATANI, Warrington Hospital, Warrington, UK; Raheel QURESHI, Queen Elizabeth Hospital, Newcastle University Medical School, Gateshead, UK; Edyta ZAGOROWICZ, The Maria Sklodowska-Curie National Research Institute of Oncology, Warsaw, Poland; Matthieu

ALLEZ, Hôpital Saint-Louis, Université Paris Nord, Paris, France; Emmanouil ARCHAVLIS, Evangelismos General Hospital, Athens, Greece; Delphine BONNET, Hôpital Rangueil, Université Toulouse 3 , Toulouse, France; Luisa GUIDI, Policlinico A. Gemelli, Università Cattolica del Sacro Cuore, Roma, Italy; Deirdre MCNAMARA, TUH, TCD, Dublin, Ireland; Piero VERNIA, Policlinico Umberto I, Università La Sapienza, Roma, Italy; Michael WEIDENHILLER, Facharztzentrum Regensburg, Regensburg, Germany; Lang ALON, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Trine BOYSEN, Hvidovre Hospital, University of Copenhagen, Hvidovre, Denmark; Charlotte DELATTRE, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Richard FARRELL, Connolly Hospital, RCSI, Dublin, Ireland; Rolf-Achim KRÜGER, Praxengemeinschaft, Radebeul, Germany; Thierry PAUPARD, CH Dunkerque, Dunkerque, France; Bénédicte PROST, CH Saint-Joseph Saint-Luc, Lyon, France; Ida VIND, Hvidovre Hospital, University of Copenhagen, Hvidovre, Denmark; Flavio CAPRIOLI, Ospedale Maggiore Policlinico di Milano, Università di Milano, Milano, Italy; Vladimir GANCHO, Nykøbing Falster Sygehus, University of Copenhagen, Nykøbing Falster, Denmark; Vincent QUENTIN, Hôpital Yves Le Foll, Saint-Brieuc, France; Benjamin AVIDAN, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Geert D'HAENS, Amsterdam University Medical Center, location AMC, Amsterdam, The Netherlands; Jane MCCARTHY, Mercy University Hospital, UCC, Cork, Ireland; Jonathon SNOOK, Poole Hospital, Poole, UK; Konstantinos SOUFLERIS, Theageneio Cancer Hospital, Thessaloniki, Greece; Frank ZERBIB, CHU Bordeaux, Université de Bordeaux, Pessac, France; Dan CARTER, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Annekatrien DEPLA, Slotervaart ziekenhuis, Amsterdam, The Netherlands; Thomas EISENBACH, Praxengemeinschaft, Leverkusen, Germany; Walter FRIES, Policlinico G. Martino, Università di Messina, Messina, Italy; Nikolaos GRAMMATIKOS, Hippokration General Hospital,

Thessaloniki, Greece; Saskia ILEGEMS, Emmaüs ziekenhuis, AZ St-Maarten, Mechelen, Belgium; Antonio LOPEZ-SANROMAN, Hospital Universitario Ramón y Cajal, University of Alcalá de Henares, Madrid, Spain; Jacques MOREAU, Hôpital Rangueil, Université Toulouse 3, Toulouse, France; Gabriele RIEGLER, Azienda ospedaliera Seconda Università di Napoli, Seconda Università di Napoli, Napoli, Italy; Svend RIETDIJK, OLVG oost, Amsterdam, The Netherlands; Marta ROCHA, Centro Hospitalar do Porto, Porto, Portugal; Isabelle ROSA, Centre hospitalier intercommunal de Créteil, Créteil, France; Barbara RYAN, TUH, TCD, Dublin, Ireland; Yelena YEREMENKO, Haemeq Medical Center, Faculty of Medicine Technion, Afula, Israel; Arnaud BORUCHOWICZ, Hôpital de Valenciennes, Université de Lille, Valenciennes, France; Filipe DAMIÃO, Centro Hospitalar Lisboa Norte, Lisboa, Portugal; Foteini LAOUDI, Athens Medical P. Faliro, Athens, Greece; Andreas LÜGERING, MVZ Portal 10, Münster, Germany; Giampiero MACARRI, Ospedale civile Augusto Murri, Fermo, Italy; Konstantinos THOMOPOULOS, University Hospital of Patra, University of Patras, Patra, Greece; Luísa BARROS, Centro Hospitalar do Tâmega e Sousa, Penafiel, Portugal; Thomas BLIXT, Zealand University Hospital, University of Copenhagen, Koege, Denmark; Aurélien GARROS, CH Saint-Joseph Saint-Luc, Lyon, France; Sam KHORRAMI, Hospital Universitario Son Espases, Palma de Mallorca, Spain; Harry SOKOL, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Andreas STURM, DRK-Kliniken Berlin, DRK-Kliniken Berlin, Berlin, Germany; Dan LIVOVSKY, Shaare Tzedek Medical Center, Hebrew University, Jerusalem, Israel; Jochen MAUL, MVZ für Gastroenterologie am Bayrischen Platz, Berlin, Germany; Heinrich MIKS, MVZ Hamm, Hamm, Germany; Vasileios PAPADOPOULOS, Attikon University Hospital, Athens, Greece; Carsten SCHMIDT, Medizinische Klinik Fulda, Medizinische Klinik Fulda, Fulda, Germany; Yifat SNIR, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Lise SVENNINGSEN, Regional Hospital Herning, University of Aarhus, Herning, Denmark;

Wafaa AHMED, Maidstone Hospital, Maidstone, UK; Yelena BROITMAN, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Emmanuel CUIILLERIER, Hôpital Victor Jousselin, Dreux, France; Prashant KANT, York Hospital, Hull York Medical School, York, UK; Jan LEYDEN, MMUH, UCD, Dublin, Ireland; Lev LICHTENSTEIN, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Susana LOPES, Centro Hospitalar Universitário de São João, Faculty of Medicine of the University of Porto, Porto, Portugal; Chloé MARTINEAU, Hôpital Saint-Antoine, Sorbonne Université, Paris, France; Hugh MULCAHY , St Vincent University Hospital & University College Dublin, Dublin, Ireland; Axel SCHWEITZER, Gastroenterologische Gemeinschaftspraxis Münster, Münster, Germany; Fiona VAN SCHAIK, UMC Utrecht, Utrecht University, Utrecht, The Netherlands; Hagar BANAI, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Pauline DANION, Hôpital Lyon-Sud, Université Lyon 1, Lyon, France; Charlotte DULERY, CHU Bordeaux, Université de Bordeaux, Pessac, France; Herma FIDDER, UMC Utrecht, Utrecht University, Utrecht, The Netherlands; Claire GAY, Hôpital Lyon-Sud ,Université Lyon 1, Lyon, France; Hervé HAGEGE, Centre hospitalier intercommunal de Créteil, Créteil, France; Florence HARNOIS, CH Marne la vallée, Jossigny, France; Søren Peter JØRGENSEN, Aarhus University Hospital, University of Aarhus, Aarhus , Denmark; Jens MÜLLER-ZIEHM, Gastroenterologische Schwerpunktpraxis Hannover, Hannover, Germany; Michail OIKONOMOU, Euroclinic, Athens, Greece; Carolina PALMELA, Hospital Beatriz Ângelo, Loures, Portugal; Jörg SCHULZE/RÖSKE, Berufsausübungsgemeinschaft, Jerichow, Germany; Mark SMITH, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Tamar THURM, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Francesca BRESSO, Karolinska University Hospital, Inst Med Solna Karolinska Institutet, Stockholm, Sweden; Hedia BRIXI,

CHU de Reims, Université de Reims Champagne Ardenne, Reims, France; John JONES, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Pdraig MACMATHUNA, MMUH, UCD, Dublin, Ireland; Spyridon MICHPOULOS, Alexandra General Hospital, Athens, Greece; Claire PAINCHART, Hôpital de Valenciennes, Université de Lille, Valenciennes, France; Yulia RON, Sourasky Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Marianne VESTER-ANDERSEN, Zealand University Hospital, University of Copenhagen, Koege, Denmark; Gonçalo ALEXANDRINO, Hospital Prof. Doutor Fernando Fonseca, Amadora, Portugal; Norbert BÖRNER, Gastroenterologische Gemeinschaftspraxis Mainz, Mainz, Germany; Mariana CARDOSO, Hospital Prof. Doutor Fernando Fonseca, Amadora, Portugal; Cristina CHAGAS, Centro Hospitalar Lisboa Ocidental, Lisboa, Portugal; Axel DIGNAß, Agaplesion Krankenhaus Frankfurt, Agaplesion Krankenhaus Frankfurt, Frankfurt, Germany; Iris DOTAN, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Charlotte HEDIN, Karolinska University Hospital, Inst Med Solna Karolinska Institutet, Stockholm, Sweden; Pantelis KARATZAS, Laiko General Hospital, Athens, Greece; Panagiotis KASAPIDIS, Central Clinic, Athens, Greece; Károly PALATKA, 2nd Department of Internal Medicine, University of Debrecen, Debrecen, Hungary; Georgios SAKIZLIS, Latriko Athinon Hospital, Athens, Greece; Ana WILSON, St Marks Hospital, Imperial college london, London, UK; Nick BOSANKO, NHS Plymouth, Plymouth, UK; Paulo CALDEIRA, Centro Hospitalar Universitário do Algarve - Faro, Faro, Portugal; Charlotte GAGNIERE, Hôpital Henri Mondor, Université de Paris, Paris, France; Louise LIBIER, Hôpital Tourcoing, Université de Lille, Tourcoing, France; Camille MEUNIER, Hôpital Lyon-Sud, Université Lyon 1, Lyon, France; Gero MOOG, Gastroenterologische Facharztpraxis, Kassel, Germany; Audrey PASQUION, CH Saint-Joseph Saint-Luc, Lyon, France; Roberta PICA, Ospedale Sandro Pertini, Roma, Italy; Ayesha AKBAR, St Marks Hospital, Imperial college london,

London, UK; Nadia ARAB, Hôpital L'archet, Nice, France; Guillaume CADIOT, CHU de Reims, Université de Reims Champagne Ardenne, Reims, France; João CARVALHO, Centro Hospitalar Vila Nova de Gaia/Espinho, Vila Nova de Gaia, Portugal; Claire CHARPIGNON, Institut Mutualiste Montsouris, Paris, France; Klaus FELLERMANN, Praxisklinik Nürnberg, Praxisklinik Nürnberg, Freudenstadt, Germany; Sigal FISHMAN, Sourasky Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Gerald FRASER, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Nathan GLUCK, Sourasky Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Mark HOESL, Praxisklinik Nürnberg, Nürnberg, Germany; Jarosław KIERKUS, Children's Memorial Health Institute, Warsaw, Poland; Maria KLOPOCKA, Nicolaus Copernicus University in Toruń, Collegium Medicum, Bydgoszcz, Poland; Eduardo MARTIN ARRANZ, Hospital Universitario La Paz, Instituto de Investigacion Sanitaria La Paz (IDIPAZ), Madrid, Spain; Luis MENCHEN, Hospital Universitario Gregorio Marañón, IiSGM, Complutense University of Madrid, Madrid, Spain; Susanna NIKOLAUS, Universitätsklinikum Schleswig-Holstein, UKSH Kiel, Kiel, Germany; Anca PETRACHE, Vejle Hospital, University of Odense, Vejle, Denmark; Cyriel PONSIOEN, Amsterdam UMC Location AMC, Amsterdam, The Netherlands; Sabino RIESTRA, Hospital Universitario Central de Asturias, Instituto de Investigación Sanitaria del Principado de Asturias (ISPA), University of Oviedo, Oviedo, Spain; Pilar ROBLEDO, Hospital Universitario de Cáceres, University of Extremadura, Cáceres, Spain; Cristina RODRIGUEZ, Complejo Hospitalario de Navarra, Public University of Navarra, Pamplona, Spain; Misheal SAMER, Haemeq Medical Center, Faculty of Medicine Technicion, Afula, Israel; Matthias TISCHER, Gastroenterologische Praxis Dresden, Dresden, Germany; Joanna WYPYCH, Copernicus Hospital, Gdańsk, Poland; Julien BAUDON, Centre Hospitalier de Cholet, Cholet, France; Cristina BEZZIO, Ospedale di Rho, Rho, Italy; Gilles BOSCHETTI, Hôpital Lyon-Sud,

Université Lyon 1, Lyon, France; Johan BURISCH, Hvidovre Hospital, University of Copenhagen, Hvidovre, Denmark; Tom CREED, Bristol Royal Infirmary, University of Bristol, Bristol, UK; Maria Giulia DEMARZO, Azienda ospedaliera universitaria San Martino, Università di Genova, Genova, Italy; Stefano FESTA, Ospedale San Filippo Neri, Roma, Italy; Andrés FIGUEROA, Hospital Universitario de Gran Canaria Dr. Negrín, University of Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain; Mette JULSGAARD, Aarhus University Hospital, University of Aarhus, Aarhus, Denmark; Pablo NAVARRO, Hospital Clinico Universitario de Valencia, University of Valencia, Valencia, Spain; Pablo PEREZ-GALINDO, Complejo Hospitalario Universitario de Pontevedra, Pontevedra, Spain; Cléa ROUILLON, CHU Caen, Université de Caen, Caen, France; Emanuele SABLICH, Ospedale di Rho, Rho, Italy; Joan TOSCA, Hospital Clinico Universitario de Valencia, University of Valencia, Valencia, Spain; Mathias VIDON, CHU Trousseau Tours, Créteil, France; Marine VIDON, CHRU Nancy, Université de Nancy, Nancy, France; René-Louis VITTE, Centre Hospitalier Poissy St Germain-en-Laye, St Germain-en-Laye, France ; Anne WAMPACH, Centre hospitalier intercommunal de Créteil, Créteil, France;

National Coordinators :

Vered ABITBOL, Hôpital Cochin AP-AH, Université Paris Cité, Paris, France; Sebastian SHAJI, Hull University Teaching Hospitals - Hull York Medical School, University of Hull, Hull, UK; Livia BIANCONE, Policlinico Tor Vergata, Università di Roma Tor Vergata, Roma, Italy; Alessandro ARMUZZI, IBD Center Humanitas Research Hospital, Rozzano Department of Biomedical Sciences Humanitas University, Peive Emanuele, Milan, Italy; Konstantinos KARMIRIS, Venizeleio General Hospital, Heraklion, Greece; Nikolaos VIAZIS, Evangelismos General Hospital, Athens, Greece; Eugeni DOMENECH, Hospital Universitari Germans Trias i Pujol Centro de Investigación Biomédica en Red de Enfermedades Hepáticas

y Digestivas (CIBEREHD), Universitat Autònoma Barcelona, Badalona, Spain; Javier P. GISBERT, Hospital Universitario de La Princesa, Instituto de Investigación Sanitaria Princesa (IIS-IP) and Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Universidad Autónoma de Madrid (UAM), Madrid, Spain; Peter BOSSUYT, Imelda General Hospital, Bonheiden, Belgium; Catherine REENAERS, CHU de Liège, ULiège, Liège, Belgium; Christian MAASER, Lüneburg Clinic, Lüneburg Clinic , Lüneburg, Germany; Britta SIEGMUND, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Ulf HELWIG, Private Practice for Internal Medicine, University of Kiel, Germany; Johan BURISCH, Copenhagen University Hospital - Amager and Hvidovre, Hvidovre, Denmark; Mette JULSGAARD, Aarhus University Hospital, University of Aarhus , Aarhus, Denmark; Henit YANAI, Rabin Medical Center, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; Uri KOPYLOV, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Glen DOHERTY, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Fernando MAGRO, Centro Hospitalar Universitário de São João, Faculdade de Medicina da Universidade do Porto, Porto, Portugal; Tamás MOLNAR, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Mark LöWENBERG, Amsterdam University Medical Center, location AMC, Amsterdam, The Netherlands; Bas OLDENBURG, UMC Utrecht, Utrecht University, Utrecht, The Netherlands; Jonas HALFVARSON, Örebro University Hospital, Faculty of Medicine and Health of Örebro University, Örebro, Sweden; Ann-Sofie BACKMAN, Karolinska University Hospital, Inst Med Solna Karolinska Institutet, Stockholm, Sweden; Edyta ZAGOROWICZ, The Maria Sklodowska-Curie National Research Institute of Oncology, Warsaw, Poland; Jarosław KIERKUS, Children's Memorial Health Institute, Warsaw, Poland;

Methodologist / Biostatistician / Data manager :

Cédric BAUMANN, UMDS CHRU Nancy - Hôpitaux de Brabois, Vandoeuvre-Les-Nancy, France; Isabelle CLERC URMES, UMDS CHRU Nancy - Hôpitaux de Brabois, Vandoeuvre-Les-Nancy, France; Hélène ROUSSEAU, UMDS CHRU Nancy - Hôpitaux de Brabois, Vandoeuvre-Les-Nancy, France; Marc BORIE, UMDS CHRU Nancy - Hôpitaux de Brabois, Vandoeuvre-Les-Nancy, France.

Coders :

Bénédicte CARON, Hôpital Universitaire de Strasbourg, Université de Strasbourg, Strasbourg, France ; Julien KIRCHGESNER, Hôpital Saint-Antoine, Sorbonne Université, Paris, France ; Mathieu UZZAN, Hôpital Henri Mondor, Université Paris Est, Créteil, France; Rachel COONEY, Queen Elizabeth Hospital, University Of Birmingham, Birmingham, UK; Kelly CHATTEN, Fairfield General Infirmary, Oldham, UK; Jimmy LIMDI, Northern Care Alliance NHS Foundation Trust, University of Manchester, Manchester, UK; Rimmer PETER, Queen Elizabeth Hospital, University Of Birmingham, Birmingham, UK; Iqbal TARIQ, Queen Elizabeth Hospital, University Of Birmingham, Birmingham, UK; Marta COSSIGNANI, Policlinico Tor Vergata, Università Di Roma Tor Vergata, Roma, Italy; Nikolaos VIAZIS, Evangelismos General Hospital, Athens Greece; Fiorella CAÑETE, Hospital Universitari Germans Trias I Pujol Centro De Investigación Biomédica En Red De Enfermedades Hepáticas Y Digestivas (CIBEREHD), Badalona, Spain ; Tom HOLVOET, Bird Group, Landegem, Belgium; Susanne KRASZ, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany; Petra WEIMERS, North Zealand Hospital, University Of Copenhagen, Hillerød, Denmark; Idan GOREN, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan_Israel, Juliette SHERIDAN, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Sandra DIAS, Grupo De Estudo Da Doença Inflamatória Intestinal - GEDII, Porto, Portugal; Klaudia FARKAS, Albert Szent-

Györgyi Health Center, University Of Szeged, Szeged, Hungary; Daniel BERGEMALM, Örebro University Hospital, Faculty of Medicine and Health, Örebro University, Örebro_Sweden; Edyta ZAGOROWICZ, The Maria Sklodowska-Curie National Research Institute of Oncology, Warsaw, Poland.

Study Coordinators:

Hadas ABALIA, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Aziza ABAZA, Hôpital de Meaux, Meaux, France; Gal ABRAMOVICH, Haemeq Medical Center, Faculty of Medicine Technion, Afula, Israel; Ingrid ACKZELL, Karolinska University Hospital, Stockholm, Sweden; Carol ADAMS, University Hospital of North Tees, Stockton, UK; Catherine ADDLETON, Royal Derby Hospitals, Derby, UK; Erika ALFAMBRA, Hospital Clínico Universitario "Lozano Blesa". Instituto de Investigación Sanitaria de Aragón (IIS-Aragón), Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Zaragoza, Spain; Alicia ALGABA, Hospital Universitario de Fuenlabrada, University Rey Juan Carlos, Madrid, Spain; Clare ALLCOCK, Russells Hall Hospital, Dudley, UK; Joanna ALLISON, Yeovil District Hospital, Yeovil, UK; Karine AMOURIAUX, CHU Nantes, Université de Nantes, Nantes, France; Julie ANDERSON, York Hospital, Hull York Medical School, York, UK; Emma ANDERSON, Our Lady of Lourdes Hospital , RCSI, Drogheda, Ireland; Saskia Appelmans, BIRD GROUP, Landegem, Belgium; Sophie ARMAND, CH Saint-Joseph Saint-Luc, Lyon, France; Lisa ARMSTRONG, Airedale General Hospital, Leeds University School of Medicine, Airedale, UK; Stacey ATKINS, Torbay Hospital, Torquay, UK; Masoumeh ATTARAN-BANDARABADI, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover,

Hannover, Germany; Yvonne BAILEY, TUH, TCD, Dublin, Ireland; Florian BÄR, Schwerpunktpraxis Gastroenterologie, Lübeck, Germany; Stephanie BARDOT, CHU Nantes, Université de Nantes, Nantes, France; Natasha BECK, Bristol Royal Infirmary, University of Bristol, Bristol, UK; Lillie BENNETT, Queen Elizabeth Hospital, University of Birmingham, Birmingham, UK; Jonathan Phil BERGFELD, Medizinische Klinik Fulda, Medizinische Klinik Fulda, Fulda, Germany; Ramdane BERKANE, Hôpital de Meaux, Meaux, France; Marion BLIN, Hôpital Nord Marseille, Université Aix-Marseille, Marseille, France; Hanne BOEY, Imelda Hospital, Bonheiden, Belgium; Louise BOWLAS, Queen Elizabeth Hospital, University of Birmingham, Birmingham, UK; Joanne BRADLEY-POTTS, Wythenshawe Hospital, University of Manchester, Manchester, UK; Tracy BREAR, King's Mill Hospital, University of Nottingham, Sutton-in-Ashfield, UK; Nicole BRETLENDER-PETERS, MVZ Hamm, Hamm, Germany; Ellen BROWN, Darlington Memorial Hospital, Durham, UK; Johanna BROWN, Royal Derby Hospitals, Derby, UK; Elizabeth BUCKINGHAM, Royal Shrewsbury Hospital, Keele University School of Medicine, Staffordshire University, Shrewsbury, UK; Katrien BUELLENS, Emmaüs ziekenhuis, AZ St-Maarten, Mechelen, Belgium; Rhian BULL, Chelsea and Westminster Hospital, Imperial College London, London, UK; Maura Burke, UCHG, NUIG, Galway, Ireland; Leighanne BURNS, University Hospital Coventry, University of Warwick, Coventry, UK; Julie BURTON, Pinderfields Hospital, Wakefield, UK; Agness BWALYA, Harrogate District Hospital, Harrogate, UK; Karine CABANAS, CHU Nantes, Université de Nantes, Nantes, France; Muriel Callaghan, Connolly Hospital, RCSI, Dublin, Ireland; Océane CAMOU, Hôpital Rangueil, Université Toulouse 3, Toulouse, France; Debbie CAMPBELL, Colchester General Hospital, Barts and The London School of Medicine and Dentistry Norwich Medical School Anglia Ruskin University, Colchester, UK; Elvira CAPOFERRO, Ospedale Sacro Cuore Don Calabria, Negrar di Valpolicella, Italy; Mandy CARNAHAN, Royal Shrewsbury Hospital, Keele University

School of medicine, Staffordshire University, Shrewsbury, UK; Cornelia CARNIO, Universitätsklinikum Erlangen, Universitätsklinikum Erlangen, Erlangen, Germany; Anne CARTER , Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Concetta CASALI CLACK, MVZ for Gastroenterology at Bayrischer Platz, Berlin, Germany; Leïla CHEDOUBA, Hôpital Saint-Louis, Université Paris Nord, Paris, France; Bessie CIPRIANO, St Bartholomew's Hospital, Queen Mary University of London, London, UK; Sophie CLAEYS, AZ St-Lucas, Ghent, Ghent, Belgium; Elizabeth CLAYTON , Pinderfields Hospital, Wakefield, UK; Manon CLOSSET, CHRU Nancy, Université de Nancy, Nancy, France; Dilek COBAN, CHU Clermont-Ferrand, Inserm U1071, M2iSH, Université Clermont Auvergne, Clermont-Ferrand, France; Sara COCOCCIA, Policlinico San Matteo, Pavia University, Pavia, Italy; Carolann COE, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Helen COLE, Royal Devon and Exeter Hospital, University of Exeter Medical School, Exeter, UK; Emilie COLLET, CHU Namur, Site Godinne, University of Namur, Namur, Belgium; Kayleigh COLLINS, Gloucester Royal Infirmary, Gloucester, UK; Isabelle COMBES, CH Annecy Genevois, Annecy, France; Emma CONNOR, Nottingham University Hospitals NHS Trust Queen's Medical Centre, Nottingham, UK; Kathryn CONSTANTIN, Watford General Hospital, Watford, UK; Susan COOKE, Chesterfield Royal Hospital, Chesterfield, UK; Nathanaëlle CORNET, CHU Angers, Angers, France; Estelle CORRIHONS, Centre Hospitalier de la Côte Basque, Bayonne, France; Pilar CORSINO, Hospital Universitario Miguel Servet, Zaragoza, Zaragoza University, Zaragoza, Spain; Rosie CORTAVILLE , Watford General Hospital, Watford, UK; Marta COSSIGNANI, Policlinico Tor Vergata, Università di Roma Tor Vergata, Roma, Italy; Donna COTTERILL, Weston General Hospital, Weston-Super-Mare, UK; Amanda COWTON, Darlington Memorial Hospital, Durham, UK; Harriet COX, York Hospital, Hull York Medical School, York, UK; Viktoria CRIPPS, Musgrove Park Hospital, Taunton, UK; Amanda CROWDER, Chesterfield

Royal Hospital, Chesterfield, UK; Tzufit CUKIER, Shaare Tzedek Medical Center, Hebrew University, Jerusalem, Israel; Amelia DANIEL, Princess Alexandra Hospital, Harlow, UK; Chris DAWE, Royal Bolton Hospital, Bolton, UK; Jose DE HAAN, Gelre ziekenhuizen, Gelre, The Netherlands; Rosanna DE LA CROIX, OLVG oost, Amsterdam, The Netherlands, Lien DECAESTEKER, AZ Delta, campus Rumbeke, Roeselare, Belgium; Evva DEJONCKHEERE, AZ Delta, campus Rumbeke, Roeselare, Belgium; Juan DELARE VILLANEGRO, Addenbrookes's Hospital, University of Cambridge, Cambridge, UK; Guillaume DELAVAL, CHU Caen, Université de Caen, Caen, France; Mariangela DELLIPONTI, Policlinico San Matteo, Pavia University, Pavia, Italy; Aude DELOMMEZ, CHU Lille, Université de Lille, Lille, France; Emilie DETRY, CHRU Nancy, Université de Nancy, Nancy, France; Melanie DHANARATNE, Ealing Hospital, Imperial college london, London, UK; Sandra DIAS, Grupo de Estudo da Doença Inflamatória Intestinal GEDII, Porto, Portugal; LAURA DIEZ GALAN, Hospital Universitario Miguel Servet, Zaragoza, Zaragoza University, Zaragoza, Spain; Marie DODEL, CHU Clermont-Ferrand, Inserm U1071, M2iSH, Université Clermont Auvergne, Clermont-Ferrand, France; Emma DOOKS, Airedale General Hospital, Leeds University School of Medicine, Airedale, UK; Joseph DU CHEYRON, Centre Hospitalier Poissy St Germain-en-Laye, St Germain-en-Laye, France ; Linda DUANE, UCHG, NUIG, Galway, Ireland; Helene Dukwen, Internistische Praxengemeinschaft Oldenburg, Kiel University, Oldenburg, Germany; Jennifer DULLING VULGO COCHRAN, MVZ for Gastroenterology at Bayrischer Platz, Berlin, Germany; Simona DYER, Scarborough Hospital, Hull York Medical School, Scarborough, UK; Harvey DYMOND, Weston General Hospital, Weston-Super-Mare, UK; Charlotte EKBLAD, Royal United Hospitals Bath, Bath, UK; Kerry ELLIOTT, Scarborough Hospital, Hull York Medical School, Scarborough, UK; Ingrid EMMERSON, South Tyneside District Hospital, South Shields, UK; Irène EUGENE-JOLCHINE, Centre Hospitalier Poissy St Germain-en-Laye, St Germain-en-Laye, France ; Lorna FLEMING ,

Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Lorna FLEMING , Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Eve FLETCHER, Royal Cornwall Hospital, University of Exeter Medical School, Truro, UK; Sarah FORD, Hull University Teaching Hospitals, Hull York Medical School/ University of Hull, Hull, UK; Greg FORSHAW, York Hospital, Hull York Medical School, York, UK; Angela FOULDS, Torbay Hospital, Torquay, UK; Caroline FRANCOIS, Centre Hospitalier de CHOLET, Cholet, France; Nicole FUGE, Klinikum Lüneburg, Klinikum Lüneburg, Lüneburg , Germany; Gal GAFNI, Haemeq Medical Center, Faculty of Medicine Technion, Afula, Israel; Miri GANON, Wolfson Medical Center, Sackler Medical School Tel Aviv University, Tel Aviv, Israel; Olga GARCIA NUÑEZ, Hospital Universitario Gregorio Marañón, Madrid, Spain; Laura GARCIA RAMIREZ, Fundación para la Investigación Biomédica del Hospital Universitario La Paz (FIBHULP), Madrid, Spain; Sophie GELDER, Queen Elizabeth Hospital, Gateshead, UK; Raimonda GETTKOWSKI, Competence Network for Intestinal Diseases e.V., Kiel, Germany; Daniela GILARDI, Istituto clinico Humanitas, Rozzano, Italy; Paolo GIUFFRIDA, Policlinico San Matteo, Pavia University, Pavia, Italy; Vincent GOBERT, Hôpital de Meaux, Meaux, France; Jo GODDEN, West Suffolk Hospital , University of Suffolk, Suffolk, UK; Nuala GODWIN, Beaumont Hospital , RCSI, Dublin, Ireland; Kay GOULDEN, Fairfield General Infirmary, Oldham, UK; Sharon GRAHAM, Ulster Hospital, Belfast, UK; Charlotte GREEN, Barnsley General Hospital, University of Sheffield, Barnsley, UK; Marie GREEN, Newcross Hospital, Wolverhampton, UK; Aboubakar GUEYE, Groupe Hospitalier Bichat-Claude Bernard, Paris, France; Tuba GULER, Schwerpunktpraxis Gastroenterologie, Lübeck, Germany; Ida GUSTAVSSON, Örebro University Hospital, Örebro, Sweden ; Helena HADJISAVVAS, Hôpital Lyon-Sud, Université Lyon 1, Lyon, France; Fiona HAMMONDS,

Royal Cornwall Hospital, University of Exeter Medical School, Truro, UK; Christina HANTZI, Agaplesion Krankenhaus Frankfurt, Agaplesion Krankenhaus Frankfurt, Frankfurt, Germany; Marion HAUKE, Universitätsmedizin Mannheim, Universitätsmedizin Mannheim, Mannheim, Germany; Julie HAYDOCK , Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Julie HAYDOCK , Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Orla HAYES, St. James's Hospital , TCD, Dublin , Ireland; Lizette HELBO NISLEV, North Zealand Hospital, University of Copenhagen, Hillerød, Denmark; Jessica HOCHSTODTER, Competence Network for Intestinal Diseases e.V., Kiel, Germany; Ashleigh HOGG, Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Ashleigh HOGG, Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Manuela HÖLBING, Medizinische Klinik Fulda, Medizinische Klinik Fulda, Fulda, Germany; Maureen HOLLAND, Macclesfield District General Hospital, Macclesfield, UK; Maartje HOLSBERGEN , Amsterdam University Medical Center, location AMC, Amsterdam, The Netherlands; Linda HOWARD, Yeovil District Hospital, Yeovil, UK; Aviya HOYDA, Shaare Tzedek Medical Center, Hebrew University, Jerusalem, Israel; Robert HULL, Royal Bolton Hospital, Bolton, UK; Jane IRISH, St Bartholomew's Hospital, Queen Mary University of London, London, UK; Wendy JACKSON, James Cook University Hospital, Durham University, Newcastle University, Middlesbrough, UK; Wendy JANSSEN, AZ Turnhout, Turnhout, Belgium; Lesley JEFFREY , Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Lesley JEFFREY, Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Sofia JOURDAN, CHU Dijon, Dijon, France; Izabela JUTROWSKA, Medical Network, Warsaw, Poland; Chava KANIEL, Shaare Tzedek Medical Center, Hebrew University, Jerusalem, Israel; Theofilos KAREZOS, EOMIFNE (Greek IBD Club), Athens, Greece; Niamh

KELLY, Bon Secours, UCC, Cork, Ireland; Jessica KELLY, Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Mary Kennedy , Connolly Hospital , RCSI, Dublin, Ireland; Una KENNEDY , St. James's Hospital, TCD, Dublin, Ireland; Joyce KIBARU, St George's Hospital, University of London, London, UK; Gemma KIRKMAN, Doncaster Royal Infirmary, Doncaster, UK; Janine KLAPROTH, Klinikum Lüneburg, Klinikum Lüneburg, Lüneburg , Germany; Corinna KNEESE, Schwerpunktpraxis Gastroenterologie, Lübeck, Germany; Andrea KOCH, Competence Network for Intestinal Diseases e.V., Kiel, Germany; Kathleen KOKKE, AZ Turnhout, Turnhout, Belgium; Martha KOPPELOW, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Susanne KRASZ, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany; Sabine KRAUSE, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Sabine KRAUSPE, Medizinische Klinik Fulda, Medizinische Klinik Fulda, Fulda, Germany; Petra KWAKKENBOS, Slotervaart ziekenhuis, Amsterdam, The Netherlands; Nunzia LABARILE, Policlinico di Bari, Università di Bari, Bari, Italy ; Hannah LANG, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover, Hannover, Germany; Marianne LASSAILLY, Hôpital Rangueil, Université Toulouse 3, Toulouse, France; Martine LECONTE, CHU Amiens, Université de Picardie Jules Verne, Amiens, France; Linda LEPCZYNSKI, Universitätsmedizin Mannheim, Universitätsmedizin Mannheim, Mannheim, Germany; Emma LEVELL, Hampshire Hospitals, University of Southampton, Basingstoke, UK; Nina LEVHAR, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Kerstin LINDHORT, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover, Hannover, Germany; Jessica LISLE, Hull University Teaching Hospitals, Hull York Medical School, University of Hull, Hull, UK; Beatriz LOPEZ CAUCE, Hospital Universitario Gregorio Marañón, Madrid, Spain; Gabriele LORENZ, Universitätsmedizin Mannheim, Universitätsmedizin Mannheim,

Mannheim, Germany; Ambra LOVATI, Ospedale Maggiore Policlinico di Milano, Università di Milano, Milano, Italy; Tracey LOWRY, Pinderfields Hospital, Wakefield, UK; Margareta LUND, Örebro University Hospital , Örebro, Sweden ;Anne LUTZ VORDERBRÜGGE, Gastroenterologische Gemeinschaftspraxis Mainz, Mainz, Germany; Suzanne MAANSSON, North Zealand Hospital, University of Copenhagen, Hillerød, Denmark; Videsheka MADAPATHAGE , Hôpital de Meaux, Meaux, France; Maelys CHEVIKOFF, Hôpital Lyon-Sud, Université Lyon 1 , Lyon, France; Alison MAGNESS, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Orla MANLEY, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Catherine MANYONI, Torbay Hospital, Torquay, UK; Ingke MARG, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Antonella MARRA, Policlinico di Bari, Università di Bari, Bari, Italy ; Carole MARTINS, Hôpital Beaujon, Université de Paris , Clichy, France; Arianna MASSELLA, Ospedale Sacro Cuore Don Calabria, Negrar di Valpolicella, Italy; Aurore MATHIAS, CHU Bordeaux, Pessac, France; Karin MENZEL, MVZ Portal 10, Münster, Germany; Danielle MERVYN, St Vincent University Hospital & University College Dublin, Dublin, Ireland; Charlotte MINSART, Erasmus Hospital, VUB-ULB, Brussels, Belgium; Sally MITCHELL, Airedale General Hospital, Leeds University School of Medicine, Airedale, UK; Kathleen MONKS, Royal Liverpool University Hospital, University of Liverpool, Liverpool John Moores University, Liverpool School of Tropical Medicine, Liverpool, UK; Mélanie MONTERO, CHRU Nancy, Université de Nancy, Nancy, France; Alson MOORE, Leicester Royal Infirmary , Leicester University, Leicester, UK; Maren MOSER, Competence Network for Intestinal Diseases e.V., Kiel, Germany; Alison MOSS, Musgrove Park Hospital, Taunton, UK; Angela MULLEN , MMUH, UCD, Dublin, Ireland; M^a Francisca MURCIANO, Hospital del Mar. Barcelona, Autonomous University of Barcelona, Barcelona, Spain; Deanna NAYLOR, Colchester General Hospital, Barts and The

London School of Medicine and Dentistry Norwich Medical School Anglia Ruskin University, Colchester, UK; Ansgar NEHUS, Praxis für Innere Medizin, Münster, Germany; Anne NICHOLSON, Basildon and Thurrock University Hospitals NHS Trust, Basildon, UK; Sarah NÖDING, Agaplesion Krankenhaus Frankfurt, Agaplesion Krankenhaus Frankfurt, Frankfurt, Germany; Sinead Nolan , Bon Secours, UCC, Cork, Cork; Janet NÖRENBURG, Medizinische Klinik Fulda, Medizinische Klinik Fulda, Fulda, Germany; Clare NORTHCOTT, NHS Plymouth, Plymouth, UK; Pauline OATES, Sunderland Royal Hospital, Sunderland, UK; Jim O'Connell, St. James's Hospital , Dublin , Ireland; Alison o'Kelly, Colchester General Hospital, Barts and The London School of Medicine and Dentistry Norwich Medical School Anglia Ruskin University, Colchester, UK; Noam ORBACH-ZINGBOIM, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Judit OROBITG, Hospital Universitari de Bellvitge, Universitat de Barcelona, Barcelona, Spain; Charlene OTIENO, Royal Derby Hospitals, Derby, UK; Charlotte OWEN, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Sarah Patch, Poole Hospital, Poole, UK; Maor PAUKER, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Renate PAULI, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Harriet PEARSON, York Hospital, Hull York Medical School, York, UK; FALGON PEGGY, Hôpital Lyon-Sud, Université Lyon 1 , Lyon, France; Séverine PETIT, CHRU Nancy, Université de Nancy, Nancy, France; Christine PETRISSANS, Centre Hospitalier de la Côte Basque, Bayonne, France; Simona PIERGALLINI, Ospedale civile Augusto Murri, Fermo, Italy; Lucy PIPPARD, Yeovil District Hospital, Yeovil, UK; Laura PITT, York Hospital, Hull York Medical School, York, UK; Gabriella PÓCSIK, Albert Szent-Györgyi Health Center, University of Szeged, Szeged, Hungary; Yoann POHER, Nord Marseille, Université d'Aix-Marseille, Marseille, France; Chloé POMES, Hôpital Rangueil, Université Toulouse 3, Toulouse, France; Lucy PRITCHARD, NHS Plymouth, Plymouth, UK;

Laura PUCHADES, Hospital Clinico Universitario de Valencia, University of Valencia, Valencia, Spain; Sheena QUAID, Ealing Hospital, Imperial college london, London, UK; Aleem RANA, St George's Hospital, University of London, London, UK; Aleem RANA, St George's Hospital, University of London, London, UK; Dana RAYNARD, Worthing Hospital, Worthing, UK; Mykla REILLY, Stepping Hill Hospital, Stockport, UK; Sonja REINERT, Klinikum Lüneburg, Klinikum Lüneburg, Lüneburg, Germany; Manuela REINKNECHT, Gastroenterologische Facharztpraxis, Kassel, Germany; Baerbel RENNER, Universitätsklinikum Schleswig-Holstein, UKSH Kiel, Kiel, Germany; Rob Reynolds, TUH, TCD, Dublin, Dublin; Giulia RIZZUTO, Ospedale V. Cervello, Palermo University, Palermo, Italy; Matthew ROBINSON, Calderdale Royal Hospital, Halifax, UK; Joke ROBRECHTS, Imelda Hospital, Bonheiden, Belgium; Eva M^a RODRIGUEZ, Hospital Universitario de La Princesa, GETECCU (Grupo Español de Trabajo En enfermedad de Crohn y Colitis Ulcerosa) and Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Autonomous University of Madrid (UAM), Madrid, Spain; Efrat ROSENBLUM, Haemeq Medical Center, Faculty of Medicine Technion, Afula, Israel; Tamlyn RUSSEL, Musgrove Park Hospital, Taunton, UK; Ibiyemi SADARE, NHS Plymouth, Plymouth, UK; Noa SALAMA, Sheba Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Toos SCHAKEL, OLVG oost, Amsterdam, The Netherlands; Anja SCHAUER, Medizinsiche Klinik Fulda, Medizinsiche Klinik Fulda, Fulda, Germany; Elisa SCHIAVONI, Policlinico A. Gemelli, Fondazione Policlinico Universitario Gemelli, Roma, Italy; Caroline SHAW, Connolly Hospital , RCSI, Dublin , Ireland; Sarah SHELTON, King's Mill Hospital, University of Nottingham, Sutton-in- Ashfield, UK; Virginie SICART, Hôpital Rangueil, Université Toulouse 3, Toulouse, France; Elodie SIOUVILLE, CHU Nantes, Université de Nantes, Nantes, France; Orla SMITH, St Vincent University Hospital & University College Dublin, Dublin , Ireland; Théo SOUDE, CHU Nantes, Université de

Nantes, Nantes, France; Sophie STEPHENSON, Bradford Teaching Hospitals, Leeds School of Medicine, Bradford, UK; Elaine STEPHENSON, Royal Victoria Infirmary, Newcastle University Medical School, Newcastle, UK; Marjan STEPPE, BIRD GROUP, Landegem, Belgium; An STERKX, OLV-Aalst, Aalst, Belgium; Jo STICKLEY, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Jo STICKLEY, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Jo STICKLEY, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Kathleen SUGRUE, Mercy University Hospital, UCC, Cork, Ireland; Natalia SWIETEC, UZ Ghent, Ghent University, Ghent, Belgium; Charlotte TASIAUX, CHU Namur, site Godinne, University of Namur, Namur, Belgium; Bhavneet THAMU, Royal Hallamshire Hospital, University of Sheffield, Sheffield, UK; Susane THOMAS, Universitätsklinikum Carl Gustav Carus, University Hospital Carl Gustav Carus, Dresden, Germany; Ogwa TOBI, Watford General Hospital, Watford, UK; Kahina TOUABI, CHRU Nancy, Université de Nancy, Nancy, France; Shifra TOVI, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Julie TREGONNING, James Cook University Hospital, Durham University, Newcastle University, Middlesbrough, UK; Laura TURCHINI, Policlinico A. Gemelli, Fondazione Policlinico Universitario Gemelli, Roma, Italy; Julia UNKHOF, Gastroenterologische Gemeinschaftspraxis Münster, Münster, Germany; Olesya UNRUH, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover, Hannover, Germany; Nurcan UZUN, Charité Campus Benjamin Franklin, Charité Berlin, Berlin, Germany; Frauke VAN AERT, UZA, Antwerp University, Antwerp, Belgium; Sandrine VANDEN BERGH, Chwapi, Tournai, Belgium; Louise VANDENBROUCKE, AZ Damiaan, Ostend, Belgium; Laura VANSTEENKISTE, BIRD GROUP, Landegem, Belgium; Shay VARDIT, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Valentin

VERGRIETE, CH Dunkerque, Dunkerque, France; Elaine WALKER, Watford General Hospital, Watford, UK; Eleanor WARNER, Maidstone Hospital, , Maidstone, UK; Olivia WATCHORN, Leicester Royal Infirmary , Leicester University, Leicester, UK; Ekaterina WATSON, Ealing Hospital, Imperial college london, London, UK; Marie-Claire WAUTHIER, CHU St-Pierre, VUB -ULB, Brussels, Belgium; Maria WEETMAN, University Hospital of North Tees, Stockton, UK; Margaret WESTON, Leicester Royal Infirmary, Leicester University, Leicester, UK; Wiebke WEST-PETROSCHKA, Universitätsklinikum Schleswig-Holstein, UKSH Kiel, Kiel, Germany; Susann WIENECKE, Klinikum Lüneburg, Klinikum Lüneburg, Lüneburg, Germany; Kerstin WIERLING, Gastroenterologische Gemeinschaftspraxis Münster, Münster, Germany; Miriam WIESTLER, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover, Hannover, Germany; Rebecca WILCOX, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Elva WILHELMSSEN, James Paget University Hospital, University of Suffolk, Great Yarmouth, UK; Angharad WILLIAMS, West Suffolk Hospital , University of Suffolk, Suffolk, UK; Georgina WILLIAMSON, Wythenshawe Hospital, University of Manchester, Manchester, UK; Deborah WILSON, University Hospital of North Tees, Stockton, UK; Kate WISTANCE, Royal Shrewsbury Hospital, Keele University School of medicine, Staffordshire University, Shrewsbury, UK; Nicolas WORTMANN, Medizinische Hochschule Hannover, Medizinische Hochschule Hannover, Hannover, Germany; Subie WURIE, University Hospital Coventry, University of Warwick, Coventry, UK; Karin YADGAR, Rabin Medical Center, Sackler Medical School Tel Aviv University, Ramat Gan, Israel; Gail YOUNG, Ulster Hospital, Belfast, UK; Megan YOUNG, Macclesfield District General Hospital, Macclesfield, UK.

GETAID contributors :

Julien AUCOUTURIER; Marie- Jo BERTIN; Hasnae BOUGRINE; Marie COISNON;
Antoine DEFRANCE; Kati GUTIERREZ; Amel HAROUZ; Laure JERBER; Aida KHLIFI;
Amina KIRATI; Nasaladjine LIWORO ; Maude LOGOLTAT; Charlotte MAILHAT;
Chancely M'BAYI; Yasmina MEDANE; Dalal MERKHOUFA ; Saouda MOHAMED
ELHAD; Bertille MONTHE; Fanny MOYON; Pascaline RABIEGA; Jennifer SEKELA;
Charlotte THILLOY.

SANOIA contributors (Digital interaction with patient):

Naima HAMAMOUCHE, Frederic PARTISOTTI

CleanWeb contributors (eCRF):

Patrick BLANDIN; Hocine MOKHTARI; Laure COUTARD